

**ALLIANCE**

Alliance Pharma plc  
Avonbridge House  
Bath Road  
Chippenham  
Wiltshire  
SN15 2BB  
www.alliancepharmaceuticals.com

*Distributor:*



**Pharmacon (Pte) Ltd**  
8 Kaki Bukit Road 2 #04-27/28  
Ruby Warehouse Complex  
Singapore 417841  
Tel: +65 6741 8500  
Fax: +65 6741 1470  
Email: enquiry@pharmacon.com.sg

**ALOCLAIR™**

**ALOCLAIR™**

# Soothes Pain Fast!

- FAST-ACTING
- FORMS PROTECTIVE LAYER OVER MOUTH ULCER
- NON-STINGING
- LASTS FOR HOURS
- SAFE FOR CHILDREN
- AVAILABLE IN GEL, SPRAY AND ORAL RINSE



Alocclair Plus Rinse  
60ml & 120ml



Alocclair Plus Gel  
8ml



Alocclair Plus Spray  
15ml

**A NEW APPROACH TO MANAGING MOUTH ULCERS**

### Aphthous Ulcers

Broad range of applications for recurrent aphthous ulcers, oral lesions including traumatic lesions caused by dental braces and ill-fitting dentures



Sores of mucous membrane of inner cheek and gum

### Mode of action

Unique key ingredients instantly coat, hydrate, and thus soothe and help healing process:

- Polyvinylpyrrolidone (PVP) is a barrier forming agent; coats the exposed nerve endings on the mucosa bringing immediate pain relief
- Sodium hyaluronate enhances tissue hydration to promote healing.
- Glycyrrhetic acid and Aloe Vera provide soothing effect

#### Before Application



Exposed nerve endings over-stimulated

#### After Application



Exposed nerve endings protected from over-stimulation

### Contains no alcohol, analgesic, or local anesthetic

- Does not sting, numb or burn
- Does not dry oral mucosa
- Physiological pH of 6
- Minimum taste disturbance

### ALOCAIR™ Product Information

**Ingredients:** Aqua, polyvinylpyrrolidone(PVP), maltodextrin, propylene glycol, aloe barbadensis, potassium sorbate, sodium benzoate, PEG-40 hydrogenated castor oil, disodium EDTA, benzalkonium chloride, aroma, sodium saccharin, glycyrrhetic acid, sodium hyaluronate, xanthan gum (Gel only), hydroxyethylcellulose (Oral Rinse only).

#### How to use:

**For Gel:** Apply 1 or 2 drops of gel so as to cover the whole mouth ulcer or lesion. Avoid direct contact of the applicator with the lesion. Do not touch the lesion with the tongue for at least 2 minutes so as to allow for the protective film to form. Close the cap after each application.

**For Spray:** When using the product for the first time take the top off and apply vaporizer. Direct the vaporizer towards the lesions inside the mouth and spray three times or as needed to cover the affected area. The vaporizer allows the product to cover the oral lesions with maximum efficiency.

**For Oral Rinse:** Fill the measuring cap provided to the 10ml indicator mark. Rinse around the mouth for at least one minute before expectorating.

Use 3 - 4 times a day, or as needed and avoid eating or drinking for at least 1 hour after application. Alocclair™ is safe if ingested.

**Cautions:** The use of Alocclair™ is contra-indicated in any patient with a known history of hypersensitivity to any of the ingredients.

Precautions and Storage instructions: Do not use if the package appears damaged. Store at room temperature. Do not refrigerate. Keep out of reach of children.